

The Entomologist's Record

To encourage the publication of concise and useful new distribution records, corrections of previously published erroneous records, misidentifications, short field notes, and current news items about entomologists, amateur and professional, entomology departments and museums, prompt publication is offered in this department.

NEW TRICHOPTERA RECORDS FROM NORTH DAKOTA¹

S.C. Harris, R.W. Katayama, R.B. Carlson²

ABSTRACT: The known ranges of *Limnephilus canadensis* Banks, *Limnephilus janus* Ross (Limnephilidae: Trichoptera), and *Ptilostomis angustipennis* (Hagen) (Phryganeidae: Trichoptera) are extended to include North Dakota.

Limnephilus canadensis Banks and *Limnephilus janus* Ross were collected in Pembina County, North Dakota, from window-pane traps situated in shelterbelts adjacent to a slough. One male of each species was collected in late July, 1975. Both species have been reported from three midwestern areas: Alberta, Saskatchewan, and Minnesota. In addition, *L. canadensis* is known to occur in Maine and *L. janus* in Colorado (Nimmo, 1971).

Ptilostomis angustipennis (Hagen) has an eastern distribution extending as far west as Michigan (Ross, 1944). Three males of this species were collected in Pembina County, North Dakota in July, 1974 from window-pane traps situated in shelterbelts adjacent to a slough.

The authors wish to acknowledge the assistance of Dr. G.B. Wiggins, Royal Ontario Museum, in verifying the identification of *L. janus*.

LITERATURE CITED

- Nimmo, A. 1971. The adult Rhyacophilidae and Limnephilidae (Trichoptera) of Alberta and eastern British Columbia and their post-glacial origin. *Quaest. Entomol.* 7: 3-234.
Ross, H.H. 1944. The caddis flies, or Trichoptera, of Illinois. *Ill. Nat. Hist. Surv. Bull.* 23: 1-326.

¹ Accepted for publication: February 23, 1978

² Department of Entomology, North Dakota State University, Fargo, North Dakota 58102